

4

Incidence Trends by Cancer Site

4.1 Men and Women Combined




Figure 4.1: Comparison of trends in average annual age-adjusted cancer incidence for all sites, men and women, 1969–2018.

Table 4.1: Comparison of trends in average annual age-adjusted cancer incidence for all sites, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	394	376.8	(335.0, 422.3)	NA	NA	—
1974-1978	493	373.1	(337.4, 411.6)	407.9	(406.3, 409.6)	0.9
1979-1983	624	422.2	(385.7, 461.4)	424.9	(423.4, 426.3)	1.0
1984-1988	842	467.6	(433.2, 504.1)	460.0	(458.6, 461.4)	1.0
1989-1993	1,021	467.5	(436.8, 499.8)	497.9	(496.5, 499.4)	0.9
1994-1998	1,312	493.7	(465.2, 523.5)	489.5	(488.1, 490.9)	1.0
1999-2003	1,591	516.8	(489.9, 544.8)	496.7	(495.4, 498.1)	1.0
2004-2008	1,850	506.6	(482.1, 531.9)	489.0	(487.7, 490.3)	1.0
2009-2013	2,169	513.5	(490.6, 537.3)	469.4	(468.1, 470.6)	1.1

(continued)

Table 4.1: *(continued)*

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
2014-2018	2,401	497.6	(476.7, 519.2)	450.9	(449.6, 452.2)	1.1

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.2: Comparison of trends in average annual age-adjusted cancer incidence for breast (women), men and women, 1969–2018.

Table 4.2: Comparison of trends in average annual age-adjusted cancer incidence for breast (women), men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	27	41.2	(26.6, 60.8)	NA	NA	—
1974-1978	42	54.0	(38.6, 73.4)	104.8	(103.7, 105.9)	0.5
1979-1983	55	61.2	(45.2, 80.9)	108.9	(107.9, 109.9)	0.6
1984-1988	96	91.0	(72.3, 112.9)	131.0	(130.0, 132.1)	0.7
1989-1993	136	103.5	(86.2, 123.4)	135.3	(134.2, 136.3)	0.8
1994-1998	218	142.2	(123.1, 163.4)	140.2	(139.2, 141.2)	1.0
1999-2003	253	133.8	(117.2, 152.1)	141.6	(140.6, 142.6)	0.9
2004-2008	285	131.7	(116.5, 148.5)	131.7	(130.8, 132.6)	1.0
2009-2013	342	144.2	(128.7, 160.9)	133.3	(132.4, 134.2)	1.1
2014-2018	340	130.8	(116.7, 146.0)	134.4	(133.4, 135.4)	1.0

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-P130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.3: Comparison of trends in average annual age-adjusted cancer incidence for colon and rectum, men and women, 1969–2018.

Table 4.3: Comparison of trends in average annual age-adjusted cancer incidence for colon and rectum, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	73	83.7	(63.7, 108.0)	NA	NA	—
1974-1978	84	77.2	(60.4, 97.4)	62.1	(61.5, 62.8)	1.2
1979-1983	105	80.8	(64.6, 99.9)	63.9	(63.3, 64.5)	1.3
1984-1988	151	98.5	(82.0, 117.4)	64.4	(63.8, 64.9)	1.5
1989-1993	179	95.5	(81.2, 111.6)	59.1	(58.6, 59.6)	1.6
1994-1998	244	104.6	(90.9, 119.7)	55.4	(54.9, 55.9)	1.9
1999-2003	301	105.9	(93.5, 119.4)	52.9	(52.5, 53.4)	2.0
2004-2008	312	89.3	(79.0, 100.5)	46.5	(46.1, 46.9)	1.9
2009-2013	371	90.9	(81.2, 101.3)	38.9	(38.5, 39.2)	2.3
2014-2018	405	87.6	(78.8, 97.1)	36.7	(36.3, 37.1)	2.4

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.4: Comparison of trends in average annual age-adjusted cancer incidence for lung and bronchus, men and women, 1969–2018.

Table 4.4: Comparison of trends in average annual age-adjusted cancer incidence for lung and bronchus, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	43	38.2	(26.6, 53.2)	NA	NA	–
1974-1978	69	54.1	(41.5, 69.3)	55.0	(54.4, 55.5)	1.0
1979-1983	96	64.6	(51.2, 80.3)	61.1	(60.5, 61.6)	1.1
1984-1988	173	97.5	(82.7, 114.1)	66.0	(65.5, 66.5)	1.5
1989-1993	174	90.0	(76.4, 105.2)	68.3	(67.8, 68.8)	1.3
1994-1998	251	102.4	(89.4, 116.8)	67.3	(66.8, 67.8)	1.5
1999-2003	256	92.1	(80.6, 104.8)	65.2	(64.7, 65.7)	1.4
2004-2008	316	97.7	(86.7, 109.8)	62.7	(62.3, 63.2)	1.6
2009-2013	336	90.3	(80.3, 101.3)	57.4	(57.0, 57.8)	1.6
2014-2018	373	83.0	(74.3, 92.4)	52.3	(51.9, 52.8)	1.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.5: Comparison of trends in average annual age-adjusted cancer incidence for prostate (men), men and women, 1969–2018.

Table 4.5: Comparison of trends in average annual age-adjusted cancer incidence for prostate (men), men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	16	40.2	(20.7, 70.6)	NA	NA	—
1974-1978	15	35.7	(19.4, 60.4)	96.5	(95.1, 97.8)	0.4
1979-1983	35	61.7	(41.3, 88.6)	106.7	(105.5, 108.0)	0.6
1984-1988	30	46.5	(29.9, 69.0)	123.7	(122.5, 124.9)	0.4
1989-1993	79	93.5	(72.5, 118.5)	196.3	(194.8, 197.7)	0.5
1994-1998	73	74.3	(56.9, 95.3)	168.7	(167.4, 169.9)	0.4
1999-2003	110	86.6	(69.4, 106.8)	178.0	(176.8, 179.2)	0.5
2004-2008	109	67.9	(54.6, 83.4)	162.2	(161.0, 163.3)	0.4
2009-2013	111	60.1	(48.3, 73.8)	130.0	(129.1, 130.9)	0.5
2014-2018	126	61.1	(49.9, 74.2)	104.1	(103.2, 105.0)	0.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.6: Comparison of trends in average annual age-adjusted cancer incidence for kidney and renal pelvis, men and women, 1969–2018.

Table 4.6: Comparison of trends in average annual age-adjusted cancer incidence for kidney and renal pelvis, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	17	14.7	(7.8, 25.1)	NA	NA	—
1974-1978	20	12.3	(7.2, 19.6)	7.9	(7.7, 8.1)	1.6
1979-1983	17	13.3	(7.1, 22.6)	8.5	(8.3, 8.7)	1.6
1984-1988	46	29.4	(20.8, 40.4)	9.8	(9.5, 10.0)	3.0
1989-1993	43	19.5	(13.7, 27.0)	10.8	(10.6, 11.0)	1.8
1994-1998	53	20.8	(15.2, 27.7)	11.5	(11.2, 11.7)	1.8
1999-2003	64	20.2	(15.3, 26.2)	12.9	(12.7, 13.1)	1.6
2004-2008	77	21.9	(16.9, 27.9)	15.0	(14.8, 15.3)	1.5
2009-2013	112	25.0	(20.3, 30.4)	15.6	(15.4, 15.8)	1.6
2014-2018	150	28.9	(24.3, 34.1)	15.9	(15.7, 16.2)	1.8

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.7: Comparison of trends in average annual age-adjusted cancer incidence for stomach, men and women, 1969–2018.

Table 4.7: Comparison of trends in average annual age-adjusted cancer incidence for stomach, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	18	21.0	(11.4, 35.4)	NA	NA	–
1974-1978	21	14.1	(8.6, 21.7)	10.6	(10.3, 10.9)	1.3
1979-1983	32	22.2	(14.4, 32.6)	10.1	(9.8, 10.3)	2.2
1984-1988	40	22.7	(15.6, 32.1)	9.0	(8.8, 9.2)	2.5
1989-1993	49	20.3	(14.7, 27.3)	8.1	(7.9, 8.3)	2.5
1994-1998	59	19.7	(14.6, 25.9)	7.3	(7.2, 7.5)	2.7
1999-2003	71	25.4	(19.5, 32.6)	6.8	(6.6, 7.0)	3.7
2004-2008	82	22.0	(17.2, 27.8)	6.5	(6.3, 6.6)	3.4
2009-2013	93	22.2	(17.6, 27.6)	6.1	(6.0, 6.3)	3.6
2014-2018	82	17.2	(13.5, 21.5)	5.7	(5.5, 5.8)	3.0

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.8: Comparison of trends in average annual age-adjusted cancer incidence for pancreas, men and women, 1969–2018.

Table 4.8: Comparison of trends in average annual age-adjusted cancer incidence for pancreas, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	13	12.8	(6.3, 23.2)	NA	NA	–
1974-1978	14	10.9	(5.6, 19.1)	11.5	(11.2, 11.8)	0.9
1979-1983	18	11.8	(6.7, 19.2)	11.4	(11.1, 11.6)	1.0
1984-1988	26	13.2	(8.4, 19.8)	11.4	(11.2, 11.7)	1.2
1989-1993	23	13.3	(8.1, 20.6)	11.0	(10.8, 11.2)	1.2
1994-1998	37	15.5	(10.6, 21.8)	10.9	(10.8, 11.2)	1.4
1999-2003	49	15.9	(11.5, 21.4)	11.2	(11.0, 11.4)	1.4
2004-2008	54	16.6	(12.2, 21.9)	12.2	(11.9, 12.4)	1.4
2009-2013	55	14.3	(10.5, 19.1)	12.6	(12.4, 12.8)	1.1
2014-2018	84	18.5	(14.5, 23.2)	12.9	(12.7, 13.1)	1.4

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.9: Comparison of trends in average annual age-adjusted cancer incidence for cervix uteri (women), men and women, 1969–2018.

Table 4.9: Comparison of trends in average annual age-adjusted cancer incidence for cervix uteri (women), men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	11	13.4	(6.6, 24.4)	NA	NA	–
1974-1978	27	36.1	(23.3, 53.5)	12.3	(11.9, 12.7)	2.9
1979-1983	41	36.9	(25.6, 51.5)	10.1	(9.8, 10.4)	3.7
1984-1988	28	18.1	(11.6, 26.9)	9.5	(9.2, 9.8)	1.9
1989-1993	30	17.5	(11.2, 26.2)	9.6	(9.3, 9.8)	1.8
1994-1998	17	8.4	(4.8, 13.6)	8.5	(8.2, 8.7)	1.0
1999-2003	19	9.5	(5.5, 15.3)	7.3	(7.1, 7.5)	1.3
2004-2008	28	11.4	(7.5, 16.7)	6.7	(6.4, 6.9)	1.7
2009-2013	25	10.1	(6.4, 15.2)	6.6	(6.4, 6.8)	1.5
2014-2018	32	12.4	(8.4, 17.8)	6.7	(6.4, 6.9)	1.9

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.10: Comparison of trends in average annual age-adjusted cancer incidence for corpus uteri (women), men and women, 1969–2018.

Table 4.10: Comparison of trends in average annual age-adjusted cancer incidence for corpus uteri (women), men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	-	-	-	NA	NA	-
1974-1978	-	-	-	34.0	(33.4, 34.6)	-
1979-1983	-	-	-	27.9	(27.4, 28.4)	-
1984-1988	6	5.7	(2.0, 12.6)	25.6	(25.2, 26.1)	0.2
1989-1993	14	11.1	(5.9, 18.9)	25.6	(25.1, 26.0)	0.4
1994-1998	16	10.5	(5.8, 17.5)	26.2	(25.8, 26.6)	0.4
1999-2003	24	13.2	(8.3, 19.9)	25.6	(25.2, 26.1)	0.5
2004-2008	39	18.0	(12.7, 24.9)	25.7	(25.3, 26.1)	0.7
2009-2013	48	19.9	(14.5, 26.7)	27.2	(26.8, 27.6)	0.7
2014-2018	40	13.7	(9.7, 18.8)	27.7	(27.3, 28.2)	0.5

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.11: Comparison of trends in average annual age-adjusted cancer incidence for urinary bladder, men and women, 1969–2018.

Table 4.11: Comparison of trends in average annual age-adjusted cancer incidence for urinary bladder, men and women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	9	9.2	(3.6, 19.2)	NA	NA	–
1974-1978	7	7.7	(2.8, 16.8)	20.5	(20.1, 20.9)	0.4
1979-1983	11	9.4	(4.5, 17.4)	21.5	(21.2, 21.8)	0.4
1984-1988	19	11.1	(6.2, 18.3)	22.6	(22.2, 22.9)	0.5
1989-1993	12	6.2	(3.1, 11.2)	22.8	(22.5, 23.1)	0.3
1994-1998	19	7.7	(4.4, 12.5)	22.9	(22.6, 23.2)	0.3
1999-2003	33	12.6	(8.4, 18.1)	23.8	(23.5, 24.1)	0.5
2004-2008	42	12.8	(9.0, 17.6)	23.9	(23.6, 24.2)	0.5
2009-2013	47	13.1	(9.4, 17.8)	23.0	(22.7, 23.3)	0.6
2014-2018	60	13.9	(10.4, 18.3)	21.9	(21.6, 22.2)	0.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.

4.2 Men Only




Figure 4.12: Comparison of trends in average annual age-adjusted cancer incidence for all sites, men, 1969–2018.

Table 4.12: Comparison of trends in average annual age-adjusted cancer incidence for all sites, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	208	438.5	(367.6, 519.0)	NA	NA	—
1974-1978	255	384.2	(333.0, 441.0)	482.1	(479.2, 484.9)	0.8
1979-1983	320	464.8	(408.5, 526.6)	509.6	(507.1, 512.2)	0.9
1984-1988	392	474.2	(422.8, 530.2)	548.1	(545.6, 550.7)	0.9
1989-1993	510	531.9	(481.9, 585.7)	620.9	(618.3, 623.4)	0.9
1994-1998	629	537.3	(491.2, 586.5)	577.7	(575.4, 580.0)	0.9
1999-2003	744	554.3	(510.5, 600.9)	581.2	(579.0, 583.5)	1.0
2004-2008	837	511.3	(473.1, 551.8)	564.4	(562.3, 566.5)	0.9
2009-2013	985	514.9	(479.5, 552.1)	519.1	(517.2, 521.0)	1.0
2014-2018	1,135	515.3	(482.8, 549.5)	482.5	(480.5, 484.4)	1.1

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.13: Comparison of trends in average annual age-adjusted cancer incidence for lung and bronchus, men, 1969–2018.

Table 4.13: Comparison of trends in average annual age-adjusted cancer incidence for lung and bronchus, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	34	62.7	(39.9, 93.9)	NA	NA	–
1974-1978	55	85.9	(63.5, 113.5)	93.2	(92.0, 94.4)	0.9
1979-1983	68	93.7	(70.8, 121.6)	97.8	(96.7, 98.8)	1.0
1984-1988	99	116.6	(93.6, 143.7)	98.5	(97.5, 99.6)	1.2
1989-1993	109	123.6	(99.9, 151.1)	95.2	(94.3, 96.2)	1.3
1994-1998	163	140.3	(117.8, 166.0)	87.2	(86.3, 88.1)	1.6
1999-2003	147	115.1	(95.7, 137.3)	80.9	(80.0, 81.7)	1.4
2004-2008	177	121.5	(102.5, 142.8)	73.4	(72.7, 74.2)	1.7
2009-2013	181	109.8	(92.9, 128.9)	65.2	(64.5, 65.9)	1.7
2014-2018	187	93.9	(79.7, 109.8)	57.5	(56.8, 58.2)	1.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.14: Comparison of trends in average annual age-adjusted cancer incidence for colon and rectum, men, 1969–2018.

Table 4.14: Comparison of trends in average annual age-adjusted cancer incidence for colon and rectum, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	36	92.4	(59.6, 136.7)	NA	NA	—
1974-1978	45	74.5	(52.7, 102.4)	72.7	(71.6, 73.8)	1.0
1979-1983	49	82.7	(58.3, 113.9)	75.9	(74.9, 76.9)	1.1
1984-1988	68	96.3	(72.8, 125.0)	78.5	(77.5, 79.4)	1.2
1989-1993	90	103.4	(81.4, 129.6)	72.3	(71.4, 73.2)	1.4
1994-1998	127	117.6	(95.8, 142.8)	66.2	(65.4, 67.0)	1.8
1999-2003	139	107.3	(88.3, 129.2)	62.1	(61.4, 62.8)	1.7
2004-2008	148	88.7	(73.7, 105.9)	53.6	(53.0, 54.3)	1.7
2009-2013	188	99.4	(84.0, 116.7)	44.0	(43.5, 44.6)	2.3
2014-2018	192	86.2	(73.4, 100.5)	41.1	(40.5, 41.7)	2.1

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.15: Comparison of trends in average annual age-adjusted cancer incidence for prostate (men), men, 1969–2018.

Table 4.15: Comparison of trends in average annual age-adjusted cancer incidence for prostate (men), men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	16	40.2	(20.7, 70.6)	NA	NA	—
1974-1978	15	35.7	(19.4, 60.4)	96.5	(95.1, 97.8)	0.4
1979-1983	35	61.7	(41.3, 88.6)	106.7	(105.5, 108.0)	0.6
1984-1988	30	46.5	(29.9, 69.0)	123.7	(122.5, 124.9)	0.4
1989-1993	79	93.5	(72.5, 118.5)	196.3	(194.8, 197.7)	0.5
1994-1998	73	74.3	(56.9, 95.3)	168.7	(167.4, 169.9)	0.4
1999-2003	110	86.6	(69.4, 106.8)	178.0	(176.8, 179.2)	0.5
2004-2008	109	67.9	(54.6, 83.4)	162.2	(161.0, 163.3)	0.4
2009-2013	111	60.1	(48.3, 73.8)	130.0	(129.1, 130.9)	0.5
2014-2018	126	61.1	(49.9, 74.2)	104.1	(103.2, 105.0)	0.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.16: Comparison of trends in average annual age-adjusted cancer incidence for stomach, men, 1969–2018.

Table 4.16: Comparison of trends in average annual age-adjusted cancer incidence for stomach, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	12	33.8	(15.0, 65.4)	NA	NA	—
1974-1978	17	21.4	(12.4, 34.6)	15.5	(15.0, 16.0)	1.4
1979-1983	25	32.8	(20.4, 50.0)	14.9	(14.5, 15.3)	2.2
1984-1988	27	33.2	(20.5, 50.9)	13.5	(13.1, 13.9)	2.5
1989-1993	37	32.3	(22.1, 45.6)	12.2	(11.8, 12.6)	2.6
1994-1998	40	28.3	(19.5, 39.7)	11.1	(10.7, 11.4)	2.6
1999-2003	38	32.8	(22.2, 46.8)	9.8	(9.6, 10.2)	3.3
2004-2008	54	32.6	(23.2, 44.6)	9.3	(9.0, 9.6)	3.5
2009-2013	56	28.6	(20.9, 38.3)	8.7	(8.4, 8.9)	3.3
2014-2018	51	22.7	(16.5, 30.4)	7.9	(7.6, 8.1)	2.9

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.17: Comparison of trends in average annual age-adjusted cancer incidence for kidney and renal pelvis, men, 1969–2018.

Table 4.17: Comparison of trends in average annual age-adjusted cancer incidence for kidney and renal pelvis, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	8	13.9	(4.9, 30.9)	NA	NA	—
1974-1978	10	12.7	(5.7, 24.6)	11.5	(11.1, 12.0)	1.1
1979-1983	9	16.7	(6.6, 34.7)	12.5	(12.2, 12.9)	1.3
1984-1988	23	31.7	(18.9, 49.9)	14.0	(13.6, 14.4)	2.3
1989-1993	22	23.4	(14.0, 36.6)	15.1	(14.8, 15.5)	1.5
1994-1998	25	22.4	(13.7, 34.8)	15.9	(15.5, 16.3)	1.4
1999-2003	43	29.8	(20.7, 41.4)	17.8	(17.4, 18.2)	1.7
2004-2008	45	27.2	(19.0, 37.7)	20.3	(19.9, 20.7)	1.3
2009-2013	61	29.1	(21.8, 38.0)	21.5	(21.1, 21.9)	1.4
2014-2018	94	39.3	(31.2, 48.8)	21.8	(21.4, 22.2)	1.8

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.18: Comparison of trends in average annual age-adjusted cancer incidence for urinary bladder, men, 1969–2018.

Table 4.18: Comparison of trends in average annual age-adjusted cancer incidence for urinary bladder, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	7	17.5	(5.3, 42.5)	NA	NA	–
1974-1978	5	9.3	(2.7, 23.1)	36.7	(35.9, 37.5)	0.3
1979-1983	8	13.1	(5.4, 26.4)	38.7	(38.0, 39.5)	0.3
1984-1988	14	15.1	(7.4, 27.3)	41.1	(40.4, 41.8)	0.4
1989-1993	9	10.2	(4.3, 20.5)	40.9	(40.2, 41.5)	0.3
1994-1998	16	14.3	(7.7, 24.4)	40.7	(40.0, 41.3)	0.4
1999-2003	26	22.3	(13.6, 34.4)	41.9	(41.3, 42.5)	0.5
2004-2008	32	21.0	(13.7, 30.7)	42.2	(41.6, 42.8)	0.5
2009-2013	33	21.4	(14.1, 31.2)	40.2	(39.6, 40.7)	0.5
2014-2018	47	24.2	(17.1, 33.5)	37.9	(37.4, 38.5)	0.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.19: Comparison of trends in average annual age-adjusted cancer incidence for pancreas, men, 1969–2018.

Table 4.19: Comparison of trends in average annual age-adjusted cancer incidence for pancreas, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	8	16.0	(5.8, 35.0)	NA	NA	—
1974-1978	7	12.3	(4.1, 28.2)	14.7	(14.2, 15.2)	0.8
1979-1983	9	10.5	(4.8, 20.1)	13.7	(13.2, 14.1)	0.8
1984-1988	13	14.6	(7.2, 26.6)	13.6	(13.2, 14.0)	1.1
1989-1993	7	7.7	(2.7, 17.1)	12.5	(12.2, 12.9)	0.6
1994-1998	16	15.4	(8.2, 26.5)	12.5	(12.2, 12.9)	1.2
1999-2003	27	21.0	(13.1, 31.9)	12.9	(12.5, 13.2)	1.6
2004-2008	29	18.6	(11.8, 27.7)	14.1	(13.7, 14.4)	1.3
2009-2013	29	15.6	(10.0, 23.0)	14.4	(14.1, 14.7)	1.1
2014-2018	49	22.5	(16.1, 30.5)	14.6	(14.3, 14.9)	1.5

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.20: Comparison of trends in average annual age-adjusted cancer incidence for liver, men, 1969–2018.

Table 4.20: Comparison of trends in average annual age-adjusted cancer incidence for liver, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	8	9.8	(4.1, 19.9)	NA	NA	—
1974-1978	10	11.6	(5.0, 22.9)	3.1	(2.9, 3.3)	3.8
1979-1983	13	13.6	(6.2, 25.9)	3.3	(3.1, 3.5)	4.2
1984-1988	11	9.0	(3.8, 18.3)	3.4	(3.2, 3.6)	2.6
1989-1993	10	8.2	(3.0, 17.8)	4.2	(4.0, 4.4)	1.9
1994-1998	12	9.8	(4.3, 19.1)	5.2	(5.0, 5.4)	1.9
1999-2003	15	11.5	(5.7, 20.7)	6.3	(6.0, 6.5)	1.8
2004-2008	22	12.7	(7.2, 20.8)	8.3	(8.1, 8.5)	1.5
2009-2013	33	14.3	(9.3, 20.9)	9.8	(9.6, 10.1)	1.5
2014-2018	36	15.4	(10.5, 21.9)	10.0	(9.7, 10.2)	1.5

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.21: Comparison of trends in average annual age-adjusted cancer incidence for esophagus, men, 1969–2018.

Table 4.21: Comparison of trends in average annual age-adjusted cancer incidence for esophagus, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	6	18.2	(5.7, 43.4)	NA	NA	–
1974-1978	5	10.3	(2.7, 27.1)	5.6	(5.3, 5.9)	1.8
1979-1983	9	15.2	(6.0, 31.5)	5.7	(5.5, 6.0)	2.6
1984-1988	8	11.7	(5.0, 23.4)	6.2	(5.9, 6.4)	1.9
1989-1993	11	13.6	(6.3, 25.6)	6.9	(6.6, 7.2)	2.0
1994-1998	14	12.4	(6.4, 21.6)	7.4	(7.1, 7.7)	1.7
1999-2003	19	16.3	(9.3, 26.4)	8.2	(8.0, 8.5)	2.0
2004-2008	15	8.2	(4.4, 13.9)	8.6	(8.3, 8.8)	1.0
2009-2013	18	7.8	(4.5, 12.8)	8.2	(8.0, 8.5)	1.0
2014-2018	32	14.4	(9.4, 21.0)	8.1	(7.8, 8.3)	1.8

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.22: Comparison of trends in average annual age-adjusted cancer incidence for nasopharynx, men, 1969–2018.

Table 4.22: Comparison of trends in average annual age-adjusted cancer incidence for nasopharynx, men, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	9	17.0	(6.0, 37.9)	NA	NA	–
1974-1978	17	20.3	(11.8, 32.6)	0.8	(0.7, 0.9)	26.8
1979-1983	6	6.5	(2.4, 14.2)	0.8	(0.7, 0.9)	8.6
1984-1988	12	13.2	(6.0, 25.3)	0.7	(0.6, 0.8)	19.2
1989-1993	21	19.1	(11.1, 30.7)	0.7	(0.6, 0.7)	29.0
1994-1998	15	13.4	(7.1, 23.2)	0.6	(0.6, 0.7)	21.7
1999-2003	17	10.3	(5.8, 16.8)	0.5	(0.5, 0.6)	19.1
2004-2008	9	4.1	(1.8, 7.8)	0.6	(0.5, 0.6)	7.2
2009-2013	18	9.8	(5.4, 16.4)	0.5	(0.5, 0.6)	18.5
2014-2018	21	7.4	(4.5, 11.5)	0.5	(0.4, 0.6)	15.4

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.

4.3 Women Only




Figure 4.23: Comparison of trends in average annual age-adjusted cancer incidence for all sites, women, 1969–2018.

Table 4.23: Comparison of trends in average annual age-adjusted cancer incidence for all sites, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	186	338.7	(286.5, 397.7)	NA	NA	—
1974-1978	238	361.2	(312.7, 415.2)	369.8	(367.7, 371.9)	1.0
1979-1983	304	383.9	(337.2, 435.3)	379.8	(378.0, 381.7)	1.0
1984-1988	450	466.6	(420.5, 516.4)	412.5	(410.6, 414.3)	1.1
1989-1993	511	422.9	(384.5, 464.1)	422.5	(420.7, 424.3)	1.0
1994-1998	683	467.5	(431.1, 506.1)	433.4	(431.6, 435.1)	1.1
1999-2003	847	498.0	(463.4, 534.5)	439.8	(438.1, 441.5)	1.1
2004-2008	1,013	513.2	(480.6, 547.3)	435.9	(434.2, 437.6)	1.2
2009-2013	1,184	523.4	(492.6, 555.6)	435.3	(433.6, 436.9)	1.2
2014-2018	1,266	493.0	(465.1, 522.0)	431.7	(429.9, 433.5)	1.1

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.24: Comparison of trends in average annual age-adjusted cancer incidence for breast (women), women, 1969–2018.

Table 4.24: Comparison of trends in average annual age-adjusted cancer incidence for breast (women), women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	27	41.2	(26.6, 60.8)	NA	NA	—
1974-1978	42	54.0	(38.6, 73.4)	104.8	(103.7, 105.9)	0.5
1979-1983	55	61.2	(45.2, 80.9)	108.9	(107.9, 109.9)	0.6
1984-1988	96	91.0	(72.3, 112.9)	131.0	(130.0, 132.1)	0.7
1989-1993	136	103.5	(86.2, 123.4)	135.3	(134.2, 136.3)	0.8
1994-1998	218	142.2	(123.1, 163.4)	140.2	(139.2, 141.2)	1.0
1999-2003	253	133.8	(117.2, 152.1)	141.6	(140.6, 142.6)	0.9
2004-2008	285	131.7	(116.5, 148.5)	131.7	(130.8, 132.6)	1.0
2009-2013	342	144.2	(128.7, 160.9)	133.3	(132.4, 134.2)	1.1
2014-2018	340	130.8	(116.7, 146.0)	134.4	(133.4, 135.4)	1.0

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-P130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.25: Comparison of trends in average annual age-adjusted cancer incidence for colon and rectum, women, 1969–2018.

Table 4.25: Comparison of trends in average annual age-adjusted cancer incidence for colon and rectum, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	37	83.1	(56.9, 117.3)	NA	NA	—
1974-1978	39	79.0	(54.9, 110.0)	55.1	(54.3, 55.9)	1.4
1979-1983	56	80.1	(59.2, 106.0)	56.0	(55.3, 56.7)	1.4
1984-1988	83	101.2	(78.8, 128.0)	54.8	(54.1, 55.5)	1.8
1989-1993	89	88.5	(70.4, 109.8)	49.9	(49.3, 50.5)	1.8
1994-1998	117	94.1	(77.1, 113.9)	47.4	(46.9, 48.0)	2.0
1999-2003	162	105.4	(89.2, 123.7)	45.7	(45.1, 46.2)	2.3
2004-2008	164	89.5	(75.8, 105.0)	40.6	(40.1, 41.1)	2.2
2009-2013	183	85.2	(72.8, 99.1)	34.5	(34.1, 35.0)	2.5
2014-2018	213	87.8	(76.0, 100.9)	32.8	(32.3, 33.3)	2.7

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.26: Comparison of trends in average annual age-adjusted cancer incidence for lung and bronchus, women, 1969–2018.

Table 4.26: Comparison of trends in average annual age-adjusted cancer incidence for lung and bronchus, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	9	15.8	(6.8, 31.1)	NA	NA	—
1974-1978	14	21.3	(11.3, 36.5)	27.5	(27.0, 28.1)	0.8
1979-1983	28	36.3	(23.2, 54.1)	35.3	(34.8, 35.9)	1.0
1984-1988	74	80.4	(62.2, 102.3)	43.4	(42.8, 44.0)	1.9
1989-1993	65	62.8	(47.8, 80.9)	49.7	(49.1, 50.3)	1.3
1994-1998	88	69.4	(55.1, 86.3)	53.4	(52.8, 54.0)	1.3
1999-2003	109	73.3	(59.8, 89.0)	54.2	(53.6, 54.8)	1.4
2004-2008	139	79.6	(66.5, 94.5)	55.0	(54.4, 55.6)	1.4
2009-2013	155	75.7	(63.7, 89.2)	51.6	(51.1, 52.2)	1.5
2014-2018	186	75.3	(64.4, 87.5)	48.5	(48.0, 49.1)	1.6

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.27: Comparison of trends in average annual age-adjusted cancer incidence for kidney and renal pelvis, women, 1969–2018.

Table 4.27: Comparison of trends in average annual age-adjusted cancer incidence for kidney and renal pelvis, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	9	15.4	(6.3, 31.1)	NA	NA	–
1974-1978	10	11.9	(5.3, 22.9)	5.1	(4.9, 5.4)	2.3
1979-1983	8	10.3	(4.1, 21.3)	5.5	(5.3, 5.8)	1.9
1984-1988	23	27.7	(16.9, 42.9)	6.6	(6.3, 6.8)	4.2
1989-1993	21	16.4	(9.6, 26.2)	7.5	(7.3, 7.7)	2.2
1994-1998	28	20.1	(13.1, 29.6)	8.0	(7.7, 8.2)	2.5
1999-2003	21	12.1	(7.4, 18.8)	8.9	(8.6, 9.1)	1.4
2004-2008	32	17.1	(11.4, 24.6)	10.6	(10.3, 10.9)	1.6
2009-2013	51	21.4	(15.7, 28.5)	10.6	(10.3, 10.8)	2.0
2014-2018	56	20.1	(15.0, 26.3)	10.7	(10.4, 11.0)	1.9

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.28: Comparison of trends in average annual age-adjusted cancer incidence for cervix uteri (women), women, 1969–2018.

Table 4.28: Comparison of trends in average annual age-adjusted cancer incidence for cervix uteri (women), women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	11	13.4	(6.6, 24.4)	NA	NA	–
1974-1978	27	36.1	(23.3, 53.5)	12.3	(11.9, 12.7)	2.9
1979-1983	41	36.9	(25.6, 51.5)	10.1	(9.8, 10.4)	3.7
1984-1988	28	18.1	(11.6, 26.9)	9.5	(9.2, 9.8)	1.9
1989-1993	30	17.5	(11.2, 26.2)	9.6	(9.3, 9.8)	1.8
1994-1998	17	8.4	(4.8, 13.6)	8.5	(8.2, 8.7)	1.0
1999-2003	19	9.5	(5.5, 15.3)	7.3	(7.1, 7.5)	1.3
2004-2008	28	11.4	(7.5, 16.7)	6.7	(6.4, 6.9)	1.7
2009-2013	25	10.1	(6.4, 15.2)	6.6	(6.4, 6.8)	1.5
2014-2018	32	12.4	(8.4, 17.8)	6.7	(6.4, 6.9)	1.9

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.29: Comparison of trends in average annual age-adjusted cancer incidence for pancreas, women, 1969–2018.

Table 4.29: Comparison of trends in average annual age-adjusted cancer incidence for pancreas, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	5	9.6	(2.8, 23.8)	NA	NA	—
1974-1978	7	9.9	(3.8, 20.8)	9.2	(8.9, 9.5)	1.1
1979-1983	9	12.8	(5.3, 25.8)	9.7	(9.4, 10.0)	1.3
1984-1988	13	12.4	(6.5, 21.4)	10.0	(9.7, 10.3)	1.2
1989-1993	16	18.3	(10.1, 30.5)	9.7	(9.4, 10.0)	1.9
1994-1998	21	16.2	(9.8, 25.1)	9.7	(9.4, 9.9)	1.7
1999-2003	22	12.5	(7.7, 19.1)	9.8	(9.5, 10.0)	1.3
2004-2008	25	14.8	(9.4, 22.1)	10.6	(10.3, 10.8)	1.4
2009-2013	26	12.9	(8.2, 19.4)	11.2	(10.9, 11.4)	1.2
2014-2018	35	14.9	(10.2, 21.0)	11.4	(11.1, 11.7)	1.3

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.30: Comparison of trends in average annual age-adjusted cancer incidence for stomach, women, 1969–2018.

Table 4.30: Comparison of trends in average annual age-adjusted cancer incidence for stomach, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	6	12.4	(3.8, 29.6)	NA	NA	–
1974-1978	-	-	-	7.2	(6.9, 7.4)	-
1979-1983	7	11.5	(3.8, 26.6)	6.7	(6.5, 7.0)	1.7
1984-1988	13	13.5	(6.7, 24.4)	5.9	(5.7, 6.2)	2.3
1989-1993	12	9.7	(4.7, 17.5)	5.2	(5.0, 5.4)	1.8
1994-1998	19	11.7	(6.8, 19.0)	4.6	(4.4, 4.8)	2.5
1999-2003	33	20.6	(13.9, 29.5)	4.5	(4.3, 4.6)	4.6
2004-2008	28	14.2	(9.3, 20.7)	4.2	(4.0, 4.3)	3.4
2009-2013	37	16.9	(11.7, 23.6)	4.1	(3.9, 4.3)	4.1
2014-2018	31	12.2	(8.2, 17.5)	3.9	(3.7, 4.1)	3.1

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.31: Comparison of trends in average annual age-adjusted cancer incidence for thyroid, women, 1969–2018.

Table 4.31: Comparison of trends in average annual age-adjusted cancer incidence for thyroid, women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	10	9.1	(4.2, 17.2)	NA	NA	—
1974-1978	8	9.2	(3.9, 18.5)	6.7	(6.4, 7.0)	1.4
1979-1983	8	5.9	(2.4, 12.3)	6.3	(6.1, 6.5)	0.9
1984-1988	11	9.0	(4.2, 16.9)	7.0	(6.8, 7.3)	1.3
1989-1993	15	8.5	(4.2, 15.4)	8.0	(7.7, 8.2)	1.1
1994-1998	15	7.7	(4.2, 12.8)	9.8	(9.5, 10.1)	0.8
1999-2003	33	14.2	(9.7, 20.2)	13.1	(12.8, 13.4)	1.1
2004-2008	20	7.7	(4.6, 12.0)	18.6	(18.3, 19.0)	0.4
2009-2013	49	20.1	(14.6, 26.9)	23.4	(23.0, 23.8)	0.9
2014-2018	60	22.4	(16.9, 29.0)	23.0	(22.6, 23.4)	1.0

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.32: Comparison of trends in average annual age-adjusted cancer incidence for corpus uteri (women), women, 1969–2018.

Table 4.32: Comparison of trends in average annual age-adjusted cancer incidence for corpus uteri (women), women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	-	-	-	NA	NA	-
1974-1978	-	-	-	34.0	(33.4, 34.6)	-
1979-1983	-	-	-	27.9	(27.4, 28.4)	-
1984-1988	6	5.7	(2.0, 12.6)	25.6	(25.2, 26.1)	0.2
1989-1993	14	11.1	(5.9, 18.9)	25.6	(25.1, 26.0)	0.4
1994-1998	16	10.5	(5.8, 17.5)	26.2	(25.8, 26.6)	0.4
1999-2003	24	13.2	(8.3, 19.9)	25.6	(25.2, 26.1)	0.5
2004-2008	39	18.0	(12.7, 24.9)	25.7	(25.3, 26.1)	0.7
2009-2013	48	19.9	(14.5, 26.7)	27.2	(26.8, 27.6)	0.7
2014-2018	40	13.7	(9.7, 18.8)	27.7	(27.3, 28.2)	0.5

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.




Figure 4.33: Comparison of trends in average annual age-adjusted cancer incidence for ovary (women), women, 1969–2018.

Table 4.33: Comparison of trends in average annual age-adjusted cancer incidence for ovary (women), women, 1969–2018.

	Alaska Native People			US White People (SEER 9)		
	Count	Rate	95% CI	Rate	95% CI	RR
1969-1973	8	11.0	(4.6, 22.0)	NA	NA	–
1974-1978	10	11.8	(4.6, 24.7)	16.4	(16.0, 16.9)	0.7
1979-1983	8	10.3	(3.9, 22.1)	16.1	(15.8, 16.5)	0.6
1984-1988	11	10.8	(5.1, 20.0)	16.7	(16.3, 17.0)	0.6
1989-1993	20	15.7	(9.4, 24.7)	16.2	(15.9, 16.6)	1.0
1994-1998	21	12.9	(7.9, 19.8)	15.2	(14.9, 15.6)	0.8
1999-2003	12	6.7	(3.3, 12.1)	15.0	(14.7, 15.3)	0.4
2004-2008	31	15.6	(10.4, 22.5)	13.7	(13.4, 14.0)	1.1
2009-2013	26	11.2	(7.1, 16.6)	13.1	(12.8, 13.4)	0.9
2014-2018	21	7.8	(4.8, 12.1)	11.4	(11.1, 11.7)	0.7

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130) standard.

See section A.2 on page 171 for information regarding statistical methods, including years of data available for US White People.